

Site Map Your Account Support About Us
Offerings: [Medscape.com](#) [Charts](#) [Mobile](#) [Logician](#) [CBSHealthwatch](#)

Medscape Medline Abstract



☐ Include in
selection list.

Document # 63 of 200 from MEDLINE

Relevance to Query:

Maximum Relevance:

The formation of competent barrier lipids in reconstructed **human epidermis** requires the presence of **vitamin C**.

J Invest Dermatol 1997 Sep;109(3):348-55 (ISSN: 0022-202X)

Ponec M; Weerheim A; Kempenaar J; Mulder A; Gooris GS; Bouwstra J; Mommaas AM [[Find other articles with these Authors](#)]

Department of Dermatology, Leiden University Medical Center, The Netherlands.

Our analysis of epidermal lipids revealed that (glucosyl)ceramide profiles in various **human skin** equivalents are different from those of native tissue. The main difference is the reduced content in **skin** equivalents of ceramides 4-7 and especially the very low content of the most polar ceramides 6 and 7, which contain hydroxylated sphingoid base and/or fatty acid. To facilitate hydroxylation, the culture medium was supplemented with **vitamins C and E**. Although in **vitamin E**-supplemented medium lipogenesis was not affected, in **vitamin C**-supplemented medium the content of glucosylceramides and of ceramides 6 and 7 was markedly increased, both in the presence and absence of serum and irrespective the substrate used (inert or natural, populated or not with fibroblasts). The improvement of the lipid profile was accompanied by a marked improvement of the barrier formation as judged from extensive production of lamellar bodies, their complete extrusion at the stratum granulosum/stratum corneum interface, and the formation of multiple broad lipid lamellar structures in the intercorneocyte space. The presence of well-ordered lipid lamellar phases was confirmed by small-angle x-ray diffraction. Some differences between native and reconstructed **epidermis**, however, were noticed. Although the long-range lipid lamellar phase was present in both the native and the reconstructed **epidermis**, the short lamellar phase was present only in native tissue. It remains to be established whether these differences can be ascribed to small differences in relative amounts of individual ceramides, to differences in fatty acid profiles, or to differences in cholesterol sulfate, pH, or calcium gradients. The results indicate the key role **vitamin C** plays in the formation of stratum corneum barrier lipids.

Major Subject Heading(s)	Minor Subject Heading(s)	CAS Registry / EC Numbers
<ul style="list-style-type: none">• Ascorbic Acid [pharmacology]• Membrane Lipids [biosynthesis]• Skin [chemistry] [metabolism] Find other articles with similar Subjects.	<ul style="list-style-type: none">• Ceramides [analysis]• Epidermis [drug effects] [ultrastructure]• Membrane Lipids [chemistry]• Scattering, Radiation• Skin [drug effects]	<ul style="list-style-type: none">• 0 (Ceramides)• 0 (Membrane Lipids)• 50-81-7 (Ascorbic Acid)

Indexing Check Tags: Human; Support, Non-U.S. Gov't

Language: English

MEDLINE Indexing Date: 199711

Publication Type: JOURNAL ARTICLE

Unique NLM Identifier: 97427915

Journal Code: M; X

[Previous Document](#)



[Next Document](#)

[Go Back To Search Results](#)


Options for View/Export/Email

selected references in ☒ format.

selected references in ☒ ☒ format.

selected references in ☒ format.

To: Subject:
Cc: Comments:
From:



[Site Map](#) [Your Account](#) [Support](#) [About Us](#)

Medscape Search Options

☒

Select a database to search, enter a search term, then click "go." [Advanced Search Forms](#)

All material on this website is protected by copyright. Copyright © 1994-2001 by Medscape Inc. All rights reserved. This website also contains material copyrighted by 3rd parties. Medscape requires 3.x browsers or better from Netscape or